AP® BIOLOGY EQUATIONS AND FORMULAS

 \overline{x} = sample mean

n =sample size

population)

o =observed results

e = expected results

 $\Sigma = \text{sum of all}$

Statistical Analysis and Probability

Mean

Standard Deviation

$$\overline{x} = \frac{1}{n} \sum_{i=1}^{n} x_i$$

$$s = \sqrt{\frac{\sum (x_i - \overline{x})^2}{n - 1}}$$

Standard Error of the Mean

Chi-Square

$$SE_{\overline{x}} = \frac{S}{\sqrt{n}}$$

$$\chi^2 = \sum \frac{(o-e)^2}{e}$$

Chi-Square Table

| p | Degrees of Freedom | | | | | | | |
|-------|--------------------|------|-------|-------|-------|-------|-------|-------|
| value | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| 0.05 | 3.84 | 5.99 | 7.81 | 9.49 | 11.07 | 12.59 | 14.07 | 15.51 |
| 0.01 | 6.63 | 9.21 | 11.34 | 13.28 | 15.09 | 16.81 | 18.48 | 20.09 |

Degrees of freedom are equal to the number of distinct possible outcomes minus one.

Laws of Probability

If A and B are mutually exclusive, then:

$$P(A \text{ or } B) = P(A) + P(B)$$

If A and B are independent, then:

$$P(A \text{ and } B) = P(A) \times P(B)$$

Hardy-Weinberg Equations

$$p^2 + 2pq + q^2 = 1$$
 $p =$ frequency of allele 1 in a population $p + q = 1$ $q =$ frequency of allele 2 in a population

Metric Prefixes

s = sample standard deviation (i.e., the sample-based estimate of the standard deviation of the

| Factor | <u>Prefix</u> | Symbol |
|-----------------|---------------|--------|
| 10 ⁹ | giga | G |
| 10^{6} | mega | M |
| 10^{3} | kilo | k |
| 10^{-1} | deci | d |
| 10^{-2} | centi | c |
| 10^{-3} | milli | m |
| 10^{-6} | micro | μ |
| 10^{-9} | nano | n |
| 10^{-12} | pico | p |
| | | |

Mode = value that occurs most frequently in a data set

Median = middle value that separates the greater and lesser halves of a data set

Mean = sum of all data points divided by number of data points

Range = value obtained by subtracting the smallest observation (sample minimum) from the greatest (sample maximum)

Rate and Growth

| Rate |
|------|
|------|

$$\frac{dY}{dt}$$

dY = amount of change

dt = change in time

Population Growth

$$\frac{dN}{dt} = B - D$$

B = birth rate

$$D = \text{death rate}$$

Exponential Growth

$$\frac{dN}{dt} = r_{\text{max}}N$$

N = population size

$$K =$$
carrying capacity

Logistic Growth

$$\frac{dN}{dt} = r_{\text{max}} N \left(\frac{K - N}{K} \right)$$

 r_{max} = maximum per capita growth rate of population

The Solute Potential of a Solution

The water potential will be equal to the solute potential of a solution in an open

container because the pressure potential of

the solution in an open container is zero.

$$\Psi_{\rm S} = -iCRT$$

i = ionization constant (1.0 for sucrose)because sucrose does not ionize in water)

C = molar concentration

Water Potential (Ψ)

 $\Psi_{\rm p}$ = pressure potential

 $\Psi_{\rm S}$ = solute potential

 $\Psi = \Psi_p + \Psi_s$

R = pressure constant(R = 0.0831 liter bars/mole K)

 $T = \text{temperature in Kelvin} (^{\circ}\text{C} + 273)$

$$\mathbf{pH} = -\log[H^+]$$

Simpson's Diversity Index

Diversity Index =
$$1 - \sum \left(\frac{n}{N}\right)^2$$

n = total number of organisms of a particular species

N = total number of organisms of all species

Surface Area and Volume

Surface Area of a Sphere

$$SA = 4\pi r^2$$

Volume of a Sphere

$$V = \frac{4}{3}\pi r^3$$

l = length

h = height

r = radius

Surface Area of a Rectangular Solid

$$SA = 2lh + 2lw + 2wh$$

Volume of a Rectangular Solid

$$V = lwh$$
 $w = width$

Surface Area of a Cylinder

$$SA = 2\pi rh + 2\pi r^2$$

$$s =$$
length of one side of a cube

SA = surface area

Surface Area of a Cube

$$SA = 6s^2$$

$$V = s^3$$

 $V = \pi r^2 h$

V = volume