

2025



AP[®] Calculus BC

Sample Student Responses and Scoring Commentary

Inside:


Free-Response Question 2

- Scoring Guidelines**
- Student Samples**
- Scoring Commentary**

Part A (BC): Graphing calculator required**Question 2****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Curve C is defined by the polar equation $r(\theta) = 2\sin^2\theta$ for $0 \leq \theta \leq \pi$. Curve C and the semicircle $r = \frac{1}{2}$ for $0 \leq \theta \leq \pi$ are shown in the xy -plane.



(Note: Your calculator should be in radian mode.)

Model Solution**Scoring**

- A** Find the rate of change of r with respect to θ at the point on curve C where $\theta = 1.3$. Show the setup for your calculations.

$$\left. \frac{dr}{d\theta} \right|_{\theta=1.3} = 1.031003$$

The rate of change of r with respect to θ at the point on curve C where $\theta = 1.3$ is 1.031.

Answer with setup

Point 1 (P1)

Scoring Notes for Part A

- An exact answer of $\left. \frac{dr}{d\theta} \right|_{\theta=1.3} = 4\sin(1.3)\cos(1.3)$ earns **P1**.
- To earn **P1**, a response must indicate differentiation of r and provide the correct answer. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point.
 - Examples of responses with correct communication include $\left. \frac{dr}{d\theta} \right|_{\theta=1.3} = 1.031$ and $r'(1.3) = 1.031$.
 - Responses with incorrect communication, such as $r'(\theta) = 1.031$, $r' = 1.031$, or $\frac{dr}{d\theta} = 1.031$, are sufficient to earn **P1**.

B

Find the area of the region that lies inside curve C but outside the graph of the polar equation $r = \frac{1}{2}$.

Show the setup for your calculations.

For $0 \leq \theta \leq \pi$, $r(\theta) = \frac{1}{2}$ for $\theta = \theta_1 = \frac{\pi}{6} = 0.523599$ and $\theta = \theta_2 = \frac{5\pi}{6} = 2.617994$.	Integrand including $(r(\theta))^2$	Point 2 (P2)
$\frac{1}{2} \int_{\theta_1}^{\theta_2} \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta$	Integrand	Point 3 (P3)
$= 2.066769$	Answer	Point 4 (P4)
The area is 2.067 (or 2.066).		

Scoring Notes for Part B

- P2** is earned for a definite integral including $(r(\theta))^2$, such as $\int_a^b \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta$ or $\int_a^b (r(\theta))^2 d\theta$, with or without the differential $d\theta$.
- P3** is earned for a definite integral (or integrals) with a correct integrand, such as $\int_a^b \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta$ or $\int_a^b (r(\theta))^2 d\theta - \int_c^d \left(\frac{1}{2}\right)^2 d\theta$, with or without the differential $d\theta$.
- The limits $\theta_1 = \frac{\pi}{6} = 0.523599$ and $\theta_2 = \frac{5\pi}{6} = 2.617994$ and the factor $\frac{1}{2}$ are assessed in **P4**, not in **P2** or **P3**.
- P4** is earned for the correct answer, with or without supporting work. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.

- Incorrect or unclear communication between the correct integral and the correct answer is treated as scratch work and is not considered in scoring. For example:

- $\int_{\pi/6}^{5\pi/6} \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta = 4.133538$ so the area is 2.067.

Note: This response earns **P2** and **P3** for the integral. It also earns **P4** for the correct answer.

- $\int_{\pi/6}^{5\pi/6} \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta = 2.067$

Note: This response earns **P2** and **P3** for the integral. It also earns **P4** for the correct answer. (In this instance, incorrect linkage is not considered in scoring.)

- **Special case:** An indefinite integral with a correct integrand does not earn **P2**, earns **P3**, and is eligible to earn **P4** with a correct answer.
- A response of $\int_{\pi/6}^{\pi/2} \left((r(\theta))^2 - \left(\frac{1}{2}\right)^2 \right) d\theta = 2.067$, using the symmetry of the region, earns **P2**, **P3**, and **P4**.

C

It can be shown that $\frac{dx}{d\theta} = 4\sin\theta\cos^2\theta - 2\sin^3\theta$ for curve C . For $0 \leq \theta \leq \frac{\pi}{2}$, find the value of θ that corresponds to the point on curve C that is farthest from the y -axis. Justify your answer.

For $0 \leq \theta \leq \frac{\pi}{2}$, the curve C is in the first quadrant. Thus, a point on the curve will be farthest away from the y -axis when the x -coordinate attains its maximum value. This will either occur when $\frac{dx}{d\theta} = 0$ or at an endpoint of the interval

$$0 \leq \theta \leq \frac{\pi}{2}.$$

$$\frac{dx}{d\theta} = 0$$

$$\Rightarrow \theta = 0.955317$$

θ	$x(\theta) = r(\theta)\cos\theta$
0	0
0.955317	0.769800
$\frac{\pi}{2}$	0

Therefore, the value of θ for which the point on the curve is farthest from the y -axis is 0.955.

Considers $\frac{dx}{d\theta} = 0$

Point 5 (P5)

Justification

Point 6 (P6)

Answer with supporting work

Point 7 (P7)

Scoring Notes for Part C

- **P5** is earned for considering $\frac{dx}{d\theta} = 0$. **P5** is not earned by just presenting $\theta = 0.955317$.

A response that discusses the sign of $\frac{dx}{d\theta}$ changing or uses the phrase “critical points of $x(\theta)$ ” also earns **P5**.

- The value $\theta = 0.955317$ might be presented as $\arccos\left(\frac{1}{\sqrt{3}}\right)$, $\arcsin\left(\sqrt{\frac{2}{3}}\right)$, or $\arctan(\sqrt{2})$.
- To earn **P6** using a candidates test, a response must make a global argument by correctly evaluating $x(\theta)$ at $\theta = 0$, $\theta = 0.955317$, and $\theta = \frac{\pi}{2}$. The evaluations must be correct to the first digit after the decimal, rounded or truncated.
- Alternate justifications:
 - $\frac{dx}{d\theta} > 0$ for $0 < \theta < 0.955$, and $\frac{dx}{d\theta} < 0$ for $0.955 < \theta < \frac{\pi}{2}$. Therefore, $\theta = 0.955$ is the location of the absolute maximum for $x(\theta)$ on the interval $0 \leq \theta \leq \frac{\pi}{2}$.
 - Because $\frac{dx}{d\theta}$ changes sign from positive to negative at $\theta = 0.955$ (this might be presented as “ $\frac{dx}{d\theta} > 0$ for $\theta < 0.955$, and $\frac{dx}{d\theta} < 0$ for $\theta > 0.955$ ”), it is the location of a relative maximum for $x(\theta)$. And because $\theta = 0.955$ is the only critical point of $x(\theta)$ in the interval $0 \leq \theta \leq \frac{\pi}{2}$, it is the location of the absolute maximum for $x(\theta)$ on the interval.
 - Because $\frac{dx}{d\theta}\Big|_{\theta=0.955} = 0$ and $\frac{d^2x}{d\theta^2}\Big|_{\theta=0.955} < 0$, $\theta = 0.955$ is the location of a relative maximum for $x(\theta)$. And because $\theta = 0.955$ is the only critical point of $x(\theta)$ in the interval $0 \leq \theta \leq \frac{\pi}{2}$, it is the location of the absolute maximum for $x(\theta)$ on the interval.
- A response that presents only a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P6** but is eligible for **P7** with the correct answer. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.

- D** A particle travels along curve C so that $\frac{d\theta}{dt} = 15$ for all times t . Find the rate at which the particle's distance from the origin changes with respect to time when the particle is at the point where $\theta = 1.3$. Show the setup for your calculations.

$$\left. \frac{dr}{dt} \right|_{\theta=1.3} = \left(\left. \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} \right) \right|_{\theta=1.3} = 1.031003 \cdot 15 = 15.465041$$

$$\frac{dr}{d\theta} \cdot \frac{d\theta}{dt}$$

Point 8 (P8)

Answer

Point 9 (P9)

The particle's distance from the origin changes at a rate of 15.465.

Scoring Notes for Part D

- To earn **P8**, $\frac{dr}{d\theta} \cdot \frac{d\theta}{dt}$ can be presented either symbolically or numerically.
- P8** might be earned in one or more steps.
- A response of $1.031 \cdot 15$ or [answer from part A] $\cdot 15$ earns both **P8** and **P9**.
- P9** is earned for the correct answer, with or without supporting work. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.

Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS A and B on this page.

PART A

$$r = 2 \sin^2 \theta$$

$$\frac{dr}{d\theta} = 4 \sin \theta \cdot \cos \theta \quad @ \quad \theta = 1.3 \approx$$

$$\frac{dr}{d\theta} \approx 4 \sin(1.3) \cdot \cos(1.3) \approx 1.031$$

PART B

$$2 \sin^2 \theta = \frac{1}{2}$$

$$\sin^2 \theta = \frac{1}{4}$$

$$\sin \theta = \frac{1}{2}$$

$$\theta = \pi/6, 5\pi/6$$

$$A = \frac{1}{2} \int_{\pi/6}^{5\pi/6} [2 \sin^2 \theta]^2 d\theta - \frac{1}{2} \int_{\pi/6}^{5\pi/6} \left[\frac{1}{2}\right]^2 d\theta$$

$$A \approx 2.06677 \text{ units}^2$$

Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS C and D on this page.

PART C

$$4 \sin \theta \cos^2 \theta - 2 \sin^3 \theta = 0 \quad @ \quad \theta = 0.95532 < \pi/2$$

~~$$\theta = 2.18628 > \pi/2$$~~

~~$$\theta = \pi > \pi/2$$~~


$$x = 2 \sin^2 \theta \cos \theta$$

critical test

$$x(0) \approx 0$$

$$x(0.95532) \approx 0.7648$$

$$x(\pi/2) \approx 0$$



By the first derivative test, the distance x from the y -axis is at a max @ $\theta = 0.95532$ b/c $\frac{dx}{d\theta}$ goes from positive to negative, and $x(0.95532)$ is max distance on critical test for $0 \leq \theta \leq \pi/2$

to negative, moving x goes from increasing

PART D

~~$$\frac{d}{dt} = \frac{dr}{dt} + r \frac{d\theta}{dt}$$~~

~~$$\frac{d}{dt} = \frac{dr}{dt} + r \frac{d\theta}{dt}$$~~

$$r = 2 \sin^2 \theta$$

$$\frac{dr}{dt} = 4 \sin \theta \cdot \cos \theta \cdot \frac{d\theta}{dt}$$

$$\frac{dr}{dt}(1.3) = 4 \sin(1.3) \cdot \cos(1.3) \cdot 15$$

$$\approx 1.031 \cdot 15$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

0165830

Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS A and B on this page.

PART A

$$r(\theta) = 2\sin^2\theta$$

$$\frac{dr}{d\theta} = 4\sin\theta\cos\theta$$

$$\frac{dr}{d\theta}$$

$$r'(\theta) = 4\sin\theta\cos\theta$$

$$r'(1.3) = 1.031$$

~~$$r'(\theta) = 4\sin\theta$$~~
~~$$r'(1.3) = 3.854$$~~

PART B

$$\frac{1}{2} \int_0^{\pi} (2\sin^2\theta)^2 d\theta = \frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} \left(\frac{1}{2}\right)^2 d\theta = \frac{5\pi}{12}$$

$$2\sin^2\theta = \frac{1}{2}$$

$$\sin^2\theta = \frac{1}{4}$$

$$\sin\theta = \frac{1}{2}$$

$$\frac{\pi}{6} \text{ to } \frac{5\pi}{6}$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS C and D on this page.

PART C

$$4 \sin \theta \cos^2 \theta - 7 \sin^3 \theta = 0$$

$$2 \sin \theta \cos^2 \theta - \sin^3 \theta = 0$$

$$2 \sin \theta \cos^2 \theta = \sin^3 \theta$$

0

$$2 \cos^2 \theta = \sin^2 \theta$$

PART D

$$\frac{d\theta}{dt} = 15$$

$$\frac{dr}{dt} = 2 \sin \theta \cos \theta \frac{d\theta}{dt} = 15$$

15.465

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

0078269



Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS A and B on this page.

PART A

$$r(\theta) = 2 \sin^2 \theta$$

$$\left. \frac{dr}{d\theta} \right|_{\theta=1.3} = 1.031$$

PART B

$$\int_{\pi/6}^{5\pi/6} ((r(\theta))^2 - (\frac{1}{2})) d\theta$$

$$\frac{1}{2} = 2 \sin^2 \theta$$

$$= 0.2065$$

$$\theta = -\frac{\pi}{6}, \frac{\pi}{6}$$

Page 6

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2 Q2

Answer QUESTION 2 PARTS C and D on this page.

PART C

$$\frac{dk}{d\theta} = 4\sin\theta \cos^2\theta - 2\sin^3\theta$$

The particle is

PART D

$$\frac{d\theta}{dt} = 15$$

$$\theta = 1.3$$

Page 7

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

0085636



Question 2

Note: Student samples are quoted verbatim and may contain spelling and grammatical errors.

Overview

NEW for 2025: The question overviews can be found in the *Chief Reader Report on Student Responses on AP Central*.

Sample: 2A

Score: 9 (1-1-1-1-1-1-1-1-1)

The response earned 9 points: 1 point in part A, 3 points in part B, 3 points in part C, and 2 points in part D.

In part A the response earned **P1** in the last line for a correct answer of $4\sin(1.3)\cos(1.3)$ along with a correct decimal approximation of that and identifying it as the derivative of r .

In part B the response earned **P2** in line 1 for presenting a definite integral including $(r(\theta))^2$. The response earned **P3** in line 1 for presenting definite integrals with correct integrands. The response earned **P4** because the boxed answer is correct.

In part C the response earned **P5** in line 1 for setting $\frac{dx}{d\theta}$ equal to 0. The response earned **P6** because the presented justification makes a correct global argument using the candidates test. The response earned **P7** because a correct answer is presented with supporting work.

In part D the response earned **P8** in line 2 because a correct product of derivatives is presented. The response earned **P9** for the answer $4\sin(1.3) \cdot \cos(1.3) \cdot 15$ along with a correct decimal approximation of that in the last line.

Sample: 2B

Score: 6 (1-1-1-0-1-0-0-1-1)

The response earned 6 points: 1 point in part A, 2 points in part B, 1 part in part C, and 2 points in part D.

In part A the response earned **P1** in line 4 for a correct answer identified as $r'(1.3)$.

In part B the response earned **P2** in line 1 for a definite integral including $(r(\theta))^2$. The response earned **P3** in line 1 for definite integrals with correct integrands. The response did not earn **P4** because the circled answer is incorrect.

In part C the response earned **P5** in line 1 for setting $\frac{dx}{d\theta}$ equal to 0. The response did not earn **P6** because no justification is presented. The response did not earn **P7** because the presented answer of 0 is incorrect.

In part D the response earned **P8** in line 2 because a correct product of derivatives is presented. The response earned **P9** in line 3 because a correct answer is presented.

Question 2 (continued)**Sample: 2C****Score: 2 (1-1-0-0-0-0-0-0)**

The response earned 2 points: 1 point in part A, 1 point in part B, 0 points in part C, and 0 points in part D.

In part A the response earned **P1** in line 2 for presenting a correct answer and identifying it as the derivative of r at $\theta = 1.3$.

In part B the response earned **P2** in line 1 for presenting a definite integral including $(r(\theta))^2$. The response did not earn **P3** because the presented integrand is incorrect as $\frac{1}{2}$ is not squared. The response did not earn **P4** because the presented answer is incorrect.

In part C the response did not earn **P5** because $\frac{dx}{d\theta} = 0$ is not considered. The response did not earn **P6** because no justification is presented. The response did not earn **P7** because no answer is presented.

In part D the response did not earn **P8** because no product of derivatives is presented. The response did not earn **P9** because no answer is presented.