

2025



AP[®] Calculus BC

Sample Student Responses and Scoring Commentary

Inside:

Free-Response Question 6

- Scoring Guidelines**
- Student Samples**
- Scoring Commentary**

Part B (BC): Graphing calculator not allowed**Question 6****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The Taylor series for a function f about $x = 4$ is given by

$$\sum_{n=1}^{\infty} \frac{(x-4)^{n+1}}{(n+1)3^n} = \frac{(x-4)^2}{2 \cdot 3} + \frac{(x-4)^3}{3 \cdot 3^2} + \frac{(x-4)^4}{4 \cdot 3^3} + \dots + \frac{(x-4)^{n+1}}{(n+1)3^n} + \dots$$
 and converges to $f(x)$ on its interval of convergence.

	Model Solution	Scoring
A	Using the ratio test, find the interval of convergence of the Taylor series for f about $x = 4$. Justify your answer.	
	$\left \frac{(x-4)^{n+2}}{(n+2)3^{n+1}} \cdot \frac{(n+1)3^n}{(x-4)^{n+1}} \right = \left \frac{(x-4)(n+1)}{3(n+2)} \right $	Sets up ratio Point 1 (P1)
	$\lim_{n \rightarrow \infty} \left \frac{(x-4)(n+1)}{3(n+2)} \right = \left \frac{x-4}{3} \right \lim_{n \rightarrow \infty} \frac{n+1}{n+2} = \left \frac{x-4}{3} \right $	Limit of ratio Point 2 (P2)
	$\left \frac{x-4}{3} \right < 1 \text{ when } -3 < x-4 < 3.$ Thus, the series converges when $1 < x < 7$.	Interior of interval of convergence Point 3 (P3)
	When $x = 1$, the series is $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} \cdot 3}{n+1}$, which converges by the alternating series test.	Considers both endpoints Point 4 (P4)
	When $x = 7$, the series is $\sum_{n=1}^{\infty} \frac{3}{n+1}$, which diverges by limit comparison to the harmonic series.	Analysis and interval of convergence Point 5 (P5)
	Therefore, the series converges for $1 \leq x < 7$.	

Scoring Notes for Part A

- **P1** is earned by presenting a correct ratio with or without absolute values. Once earned, **P1** is banked (i.e., subsequent errors in simplification or evaluation do not impact scoring for **P1**). **P2** is not earned if there are any errors in simplification or evaluation of the limit.
- **P1** is earned for ratios mathematically equivalent to any of the following:

$$\frac{(x-4)^{n+2}}{(n+2)3^{n+1}} \cdot \frac{(n+1)3^n}{(x-4)^{n+1}}, \frac{(x-4)^{n+2}}{(n+2)3^{n+1}}, \frac{(x-4)^{n+1}}{(n+1)3^n} \cdot \frac{n3^{n-1}}{(x-4)^n}, \text{ or } \frac{(x-4)^{n+1}}{(n+1)3^n} \cdot \frac{n3^{n-1}}{(x-4)^n}.$$

- **P1** is also earned for ratios mathematically equivalent to the following reciprocal ratios:

$$\frac{(x-4)^{n+1}}{(n+1)3^n} \cdot \frac{(n+2)3^{n+1}}{(x-4)^{n+2}}, \frac{(x-4)^{n+1}}{(n+1)3^n}, \frac{(x-4)^n}{n3^{n-1}} \cdot \frac{(n+1)3^n}{(x-4)^{n+1}}, \text{ or } \frac{(x-4)^n}{n3^{n-1}} \cdot \frac{(n+1)3^n}{(x-4)^{n+1}}.$$

A response that presents any of these reciprocal ratios earns **P1** and is eligible for **P2**, **P3**, and **P4**, but not **P5**.

- A response that does not present a ratio is not eligible for **P2**.
- A response that does not use the absolute value of a ratio can earn both **P1** and **P2**.
- A response that does not use absolute value in computing the limit is eligible for **P3** if the expression $-1 < \frac{x-4}{3} < 1$ or an equivalent inequality is presented.
- A response that presents an incorrect limit of form $\frac{|x-4|}{b}$, where $b > 0$, is eligible for **P3**.

P3 is then earned for correctly finding an interval with interior $(1, 7)$ or $(4-b, 4+b)$.

Note: $|x-4| < b$ is not sufficient to earn **P3**. $x-4 < b$ can earn **P3** if this is resolved to $-b+4 < x < b+4$.

- **P4** is earned for considering both endpoints of the correct interval or both endpoints of an incorrect interval that has earned **P3**.
- To earn **P5**, a response must correctly analyze the series at $x = 1$ and $x = 7$, and present the correct interval of convergence. Naming of an appropriate test is sufficient for the analysis at each endpoint. In addition to the tests listed in the model solution, the direct comparison test to an appropriate series or the integral test may also be used.

- B** Find the first three nonzero terms and the general term of the Taylor series for f' , the derivative of f , about $x = 4$.

$$f'(x) = \frac{(x-4)}{3} + \frac{(x-4)^2}{9} + \frac{(x-4)^3}{27} + \cdots + \frac{(x-4)^n}{3^n} + \cdots$$

First three terms	Point 6 (P6)
General term	Point 7 (P7)

Scoring Notes for Part B

- **P6** is earned by presenting the first three nonzero terms in a list or as part of a polynomial or series.
- **P7** is earned by identifying the correct general term (either individually or as part of a polynomial or series).

- C** The Taylor series for f' described in part B is a geometric series. For all x in the interval of convergence of the Taylor series for f' , show that $f'(x) = \frac{x-4}{7-x}$.

The Taylor series for f' is a geometric series with first term

$$\frac{x-4}{3} \text{ and common ratio } \frac{x-4}{3}.$$

$$f'(x) = \frac{\frac{x-4}{3}}{1 - \frac{x-4}{3}} = \frac{x-4}{3 - (x-4)} = \frac{x-4}{7-x}$$

Verification **Point 8 (P8)**

Scoring Notes for Part C

- A response of $f'(x) = \frac{\frac{x-4}{3}}{1 - \frac{x-4}{3}}$ is sufficient to earn **P8**.

- D** It is known that the radius of convergence of the Taylor series for f about $x = 4$ is the same as the radius of convergence of the Taylor series for f' about $x = 4$. Does the Taylor series for f' described in part B converge to $f'(x) = \frac{x-4}{7-x}$ at $x = 8$? Give a reason for your answer.

It follows from the work in part A that the interior of the interval of convergence of the Taylor series for f' is $1 < x < 7$.

Therefore, $x = 8$ would be outside the interval of convergence of f' , and the Taylor series for f' would not converge to

$$f'(x) = \frac{x-4}{7-x} \text{ at } x = 8.$$

Answer with reason **Point 9 (P9)**

Scoring Notes for Part D

- A response of “no, $x = 8$ is outside the interval of convergence” is sufficient to earn **P9**.
- **P9** can be earned with a response consistent with an incorrect interval of convergence imported from part A.
- Alternate solutions:
 - Because the series for $f'(x)$ is geometric, this converges to $f'(x)$ for all values of x such that the common ratio $\frac{x-4}{3}$ is between -1 and 1 .

$$-1 < \frac{x-4}{3} < 1 \Rightarrow -3 < x-4 < 3 \Rightarrow 1 < x < 7$$
 - Because $x = 8$ is outside the interval $1 < x < 7$, the series for f' does not converge to $f'(x) = \frac{x-4}{7-x}$ at $x = 8$.
 - For $x = 8$, the series is given by $\sum_{n=1}^{\infty} \left(\frac{4}{3}\right)^n$, which is a geometric series with $r = \frac{4}{3} > 1$.
Therefore, the series diverges for $x = 8$.

Q6

NO CALCULATOR ALLOWED

Q6

Answer QUESTION 6 PARTS A and B on this page.

PART A

$$\sum_{n=1}^{\infty} \frac{(x-4)^{n+1}}{(n+1)3^n}$$

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$$

$$\lim_{n \rightarrow \infty} \left| \frac{(x-4)^{n+2}}{(n+2)3^{n+1}} \cdot \frac{3^n(n+1)}{(x-4)^{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{x-4}{3} \cdot \frac{(n+1)}{(n+2)} \right|$$

$$\left| \frac{x-4}{3} \right| < 1$$

$$-1 < \frac{x-4}{3} < 1$$

$$-3 < x-4 < 3$$

$$\boxed{1 < x < 7}$$

$$x=1: \sum_{n=1}^{\infty} \frac{(-3)^{n+1}}{(n+1)3^n} = \sum_{n=1}^{\infty} \frac{(-3)^{n+1}}{n+1}$$

$$\lim_{n \rightarrow \infty} \frac{3}{n+1} = 1 \quad 0 < 1 < \infty \quad \text{diverges by LCT to } \sum_{n=10}^{\infty} \frac{3}{n}$$

AST: $\lim_{n \rightarrow \infty} \frac{3}{n+1} = 0$, $\frac{3}{n+1} > \frac{3}{n+2}$ ✓
 Converges conditionally by AST

$$x=7: \sum_{n=1}^{\infty} \frac{3^{n+1}}{(n+1)3^n} = \sum_{n=1}^{\infty} \frac{3}{n+1}$$

Diverges by LCT

PART B

$$f = \frac{(x-4)^2}{6} + \frac{(x-4)^3}{27} + \frac{(x-4)^4}{4 \cdot 27}$$

$$f' = 2 \frac{(x-4)}{6} + 3 \frac{(x-4)^2}{27} + \frac{4(x-4)^3}{4 \cdot 27} + \dots$$

$$f' = \frac{x-4}{3} + \frac{(x-4)^2}{3^2} + \frac{(x-4)^3}{3^3} + \dots + \frac{(x-4)^n}{3^n}$$

$$\sum_{n=1}^{\infty} \frac{(x-4)^n}{3^n}$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

0002795



Q6

NO CALCULATOR ALLOWED

Q6

Answer QUESTION 6 PARTS C and D on this page.

PART C

Geometric series $\sum_{n=1}^{\infty} \frac{(x-4)^n}{3^n}$
 in the interval of convergence, $|r| < 1$ and the series converges to $\frac{a}{1-r}$.

$$r = \frac{x-4}{3}, \text{ and } a = a_1 = \frac{x-4}{3}$$

$$\frac{a}{1-r} = \frac{\frac{x-4}{3}}{1 - \frac{x-4}{3}} = \frac{\frac{x-4}{3}}{\frac{3-x+4}{3}} = \frac{x-4}{7-x}$$

PART D

No. The interval of convergence for f about $x=4$ is $1 \leq x < 7$, and $x=8$ is outside of that range. Since the radius of convergence is the same for f' and f , and both are centered about $x=4$, 8 must be out of the interval of convergence for f' as well.

Therefore, it must diverge and cannot converge to

$$f'(x) = \frac{x-4}{7-x} \text{ at } x=8.$$

check: geometric,
 $|r| < 1 \implies -1 < \frac{x-4}{3} < 1$
 $-1 < x < 7$
 geometric, so not
 $-1 \leq x < 7$

Q6

NO CALCULATOR ALLOWED

Q6

Answer QUESTION 6 PARTS A and B on this page.

PART A

$$\lim_{n \rightarrow \infty} \left| \frac{(x-4)^{n+2}}{(n+2)3^{n+1}} \cdot \frac{(n-1)3^n}{(x-4)^{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x-4)^{n+1} (n-1)}{(n+2)3} \right| = |x-4| \lim_{n \rightarrow \infty} \left| \frac{n-1}{3(n+2)} \right|$$

due to ratio test.

$$|x-4| \cdot \frac{1}{3} < 1, \quad -3 < x-4 < 3, \quad \boxed{1 < x < 7}$$

PART B

$$f' \approx \frac{2(x-4)}{2 \cdot 3} + \frac{3(x-4)^2}{3 \cdot 3^2} + \frac{4(x-4)^3}{4 \cdot 3^3} + \dots + \frac{(n+1)(x-4)^n}{(n+1)3^n} + \dots$$

$$f' \approx \frac{(x-4)}{3} + \frac{(x-4)^2}{3^2} + \frac{(x-4)^3}{3^3} + \dots + \frac{n(x-4)^n}{3^n} + \dots$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Q6

NO CALCULATOR ALLOWED

Q6

Answer QUESTION 6 PARTS C and D on this page.

PART C

Since f' is Geometric series $(S = \frac{a_1}{1-r})$. $\left(\left|\frac{x-4}{3}\right| < 1 \text{ at } x=4 \text{ converges}\right)$

$$a = \frac{(x-4)}{3} \quad r = \frac{x-4}{3}$$

$$S = \frac{\left(\frac{(x-4)}{3}\right)}{\left(\frac{3-x+4}{3}\right)} = \frac{x-4}{7-x} //$$

$$\frac{\frac{x-4}{3}}{1 - \left(\frac{x-4}{3}\right)}$$

$\therefore f'(x) = \frac{x-4}{7-x} //$

PART D

$f'(x) = \frac{x-4}{7-x}$ @ $x=8$?

No. Since $f'(x) = \frac{x-4}{7-x}$ is derived from where $x=4$ (center is 4) the $T(x)$ form is $\frac{(x-4)^n}{3^n}$. However, since the center is shifted to $x=8$, in $T(x)$, $(x-8)$ is needed to be used instead of $(x-4)$. Leading to different $f'(x)$ formula.

convergence

Q6

 NO CALCULATOR ALLOWED

Q6

Answer QUESTION 6 PARTS A and B on this page.

PART A

$$\lim_{x \rightarrow \infty} \left| \frac{(x-4)^{n+2}}{(n+2)3^{n+1}} \cdot \frac{(n+1)3^n}{(x-4)^{n+2}} \right|$$

$$\lim_{n \rightarrow \infty} |x-4| \quad \begin{matrix} -1 < x-4 < 1 \\ +4 & +4 & +4 \end{matrix}$$

$x=3$

$3 < x < 5$

$(3, 5)$

$\lim_{n \rightarrow \infty} \frac{(-1)^{n+1}}{(n+1)3^n}$ Converges

PART B

$$f'(x) = \frac{(x-4)^2}{12} + \frac{(x-4)^3}{81} + \frac{(x-4)^4}{16 \cdot 3^3} + \dots + \frac{(x-4)^{n+1}}{(2(n+1))3^n}$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

0070656



Q6

NO CALCULATOR ALLOWED


Q6

Answer QUESTION 6 PARTS C and D on this page.

PART C

$$\lim_{x \rightarrow \infty} \frac{(x-4)^{n+2}}{(2n+4)3^{n+1}} \cdot \frac{(2n+2)3^n}{(x-4)^{n+1}}$$

$$\lim_{x \rightarrow \infty} \frac{(x-4)^{n+2}}{(2n+4)3^{n+1}} \cdot \frac{(2n+2)3^n}{(x-4)^{n+1}}$$



PART D

No, since $x=8$ is not in the interval of convergence, which is $(3,5)$, the Taylor series cannot converge at that value.

Question 6

Note: Student samples are quoted verbatim and may contain spelling and grammatical errors.

Overview

NEW for 2025: The question overviews can be found in the *Chief Reader Report on Student Responses on AP Central*.

Sample: 6A

Score: 9 (1-1-1-1-1-1-1-1-1)

The response earned 9 points: 5 points in part A, 2 points in part B, 1 point in part C, and 1 point in part D.

In part A the response earned **P1** with the presentation of the correct ratio on the left side of line 2. The response earned **P2** with the result of the limit, $\left| \frac{x-4}{3} \right|$, in line 3. The response earned **P3** with the presentation of the interval $1 \leq x < 7$ in line 6 that has the correct interior $1 < x < 7$. The response earned **P4** with the first two infinite sums on the right after the expressions $x = 1$ and $x = 7$. The response earned **P5** with the correct analysis of the endpoints $x = 1$ and $x = 7$. For $x = 1$, the statement “converges conditionally by AST” and for $x = 7$ the statement “diverges by LCT to $\sum_{n=1}^{\infty} \frac{3}{n}$,” demonstrates that $x = 1$ is included in the interval of convergence but that $x = 7$ is not included. The correct interval is presented in the boxed statement on the left.

In part B the response earned **P6** with the correct presentation of the three terms in line 2. It is not necessary to simplify the expression, but the response is correctly simplified in line 3. The response earned **P7** with the correct presentation of the general term as part of the series expression on the right.

In part C the response earned **P8** with the verification in line 5.

In part D the response earned **P9** in lines 1 and 2 with the statement “No. The interval of convergence for f about $x = 4$ is $1 \leq x < 7$, and $x = 8$ is outside of that range.”

Sample: 6B

Score: 6 (1-1-1-0-0-1-1-1-0)

The response earned 6 points: 3 points in part A, 2 points in part B, 1 point in part C, and 0 points in part D.

In part A the response earned **P1** with the presentation of the correct ratio in line 1 on the left. The response earned **P2** with the correct limit of the ratio in line 2 on the left. The response earned **P3** with the presentation of the correct interior of convergence $1 < x < 7$ in the box. The response did not earn **P4** because the endpoints are not checked. The response did not earn **P5** because **P4** is not earned.

In part B the response earned **P6** with the correct presentation of the three terms in line 1. It is not necessary to simplify the expression, but it is correctly simplified and presented in line 2. The response earned **P7** with the correct presentation of the general term in line 1. It is not necessary to simplify the expression, but it is correctly simplified and presented in line 2.

In part C the response earned **P8** in line 2 on the right after $S =$.

In part D the response did not earn **P9** because the stated reason for the answer “No” is based on shifting the center of the interval, which is incorrect.

Question 6 (continued)**Sample: 6C****Score: 3 (1-0-1-0-0-0-0-1)**

The response earned 3 points: 2 points in part A, 0 points in part B, 0 points in part C, and 1 point in part D.

In part A the response earned **P1** with the presentation of the correct ratio on the left side of line 1. The response did not earn **P2** because an incorrect limit of $|x - 4|$ is computed. The response earned **P3** with the presentation of the correct interior $3 < x < 5$ based on the identification of $|x - 4|$ as the limit. The response did not earn **P4** because there was no attempt to check the endpoints. Note that the work in the lower left corner is crossed out. The response did not earn **P5** because **P4** was not earned.

In part B the response did not earn **P6** because the three terms presented in line 1 are incorrect. The response did not earn **P7** because the general term presented in line 2 is incorrect.

In part C the response did not earn **P8** because the correct expression is not used.

In part D the response earned **P9** with the statement “No, since $x = 8$ is not in the interval of convergence, which is $(3, 5)$, the Taylor series cannot converge at that value.” This answer is consistent with the interval found in part A.