

2025



AP[®] Chemistry

Sample Student Responses and Scoring Commentary

Inside:

Free-Response Question 5

- Scoring Guidelines**
- Student Samples**
- Scoring Commentary**

Question 5: Short Answer**4 points**

A For the correct answer: **Point 01***Tetrahedral*

B For the correct answer and a valid justification: **Point 02**

Agree. Compound Y has a larger, more polarizable electron cloud because the Si atom has more occupied electron shells than the C atom, giving compound Y stronger London dispersion forces and a higher boiling point than compound X.

C For the correct answer and a valid justification. **Point 03**

Examples of acceptable responses may include the following:

- Compound X. Compound X has weaker intermolecular forces than compound Y, so molecules of X are more likely to be in the gas phase at 82°C and would therefore have a higher vapor pressure.*
- Compound X. At 82°C, compound X has reached its boiling point, but compound Y has not. Therefore, the proportion of X molecules in the vapor phase would be much greater than that of compound Y, giving compound X the higher vapor pressure.*

D For the correct calculated value: **Point 04**

$$PV = nRT$$

$$n = \frac{PV}{RT} = \frac{(2.30 \text{ atm})(12.5 \text{ L})}{(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{K}\cdot\text{mol}})(471\text{K})} = 0.744 \text{ mol}$$

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Part A

Question 5

tetrahedral because there are 4 electron domains around Si

Part B

Yes, Si has more electron shells than C , so it has a more polarizable electron cloud than C , so compound Y must be more polarizable than compound X, so compound Y will have greater London dispersion forces and a higher boiling point than compound X.

Part C

Compound X will have the higher vapor pressure because more of compound X will be in vapor form at 82°C than compound Y because compound X's boiling point is equal to 82°C and compound Y's boiling point is greater than 82°C (98°C).

Page 12

Question 5 is continued on the next page.

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Part D

Question 5

$$PV = nRT$$

$$(2.30 \text{ atm})(12.5 \text{ L}) = n (0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1})(471.15 \text{ K})$$

$$n = 0.744 \text{ moles}$$

Page 13

Continue to Question 6.

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Question 5**Part A**

Tetrahedral

Part B

Yes that is correct. Since both compounds have the same types of IMFs, the higher boiling point goes to the compound w/ the larger LDF which is directly related to the larger molar mass.

Part C

Compound X will have a higher vapor pressure when heated b/c it has weaker IMF forces leading to a lower boiling point & therefore needing less energy to vaporize thus causing the higher vapor pressure.

Question 5 is continued on the next page.



Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Part D

Question 5

$$\frac{PV}{RT} = n$$

$$n = \frac{(2.30 \text{ atm})(12.5 \text{ L})}{(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(471 \text{ K})}$$

$$= .744 \text{ mols}$$

Page 13

Continue to Question 6.

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Part A

Question 5

The geometry around the atom or Si in compound Y is mostly made of 120° angles

Part B

I agree because the London dispersion forces will cause a higher boiling point the stronger they are.

Part C

Compound X will have the greater vapor pressure because it will have reached its boiling point, while Compound Y has not yet.

Page 12'

Question 5 is continued on the next page.

Use a pencil or a pen with black or dark blue ink. Do NOT write your name. Do NOT write outside the box.

Part D

Question 5

$$74.1 \text{ g/mol} + 40.2 \text{ g/mol}$$

When the mixture is both gases, the moles of gas particles will be 164.3 mols, because they both have become gases

[Handwritten signature]

Continue to Question 6.

Question 5

Note: Student samples are quoted verbatim and may contain spelling and grammatical errors.

Overview

NEW for 2025: The question overviews can be found in the *Chief Reader Report on Student Responses on AP Central*.

Sample: 5A

Score: 4

Point 01: 1

Part A: The point was earned for correctly identifying the geometry around the Si atom as tetrahedral.

Point 02: 1

Part B: The point was earned for stating “Yes” followed by a valid justification that “Si has more electron shells than C, so it has a more polarizable electron cloud than C, so compound Y must be more polarizable than compound X.”

Point 03: 1

Part C: The point was earned for correctly choosing compound X along with a valid justification that “more of Compound X will be in vapor form at 82°C than Compound Y” along with a valid comparison of the boiling points: “Compound Y’s boiling point is greater than 82°C.”

Point 04: 1

Part D: The point was earned for correctly calculating the number of moles and showing the supporting setup.

Sample: 5B

Score: 3

Point 01: 1

Part A: The point was earned for correctly identifying the geometry around the Si atom as tetrahedral.

Point 02: 0

Part B: The point was not earned because the response attributes the larger London dispersion forces to the “larger molar mass” rather than the larger, more polarizable electron cloud.

Point 03: 1

Part C: The point was earned for correctly choosing compound X along with a valid justification that compound X has “weaker IMF forces” leading to a lower boiling point and higher vapor pressure. The justification is valid for all temperatures, including 82°C.

Point 04: 1

Part D: The point was earned for correctly calculating the number of moles and showing the supporting setup. The incorrect value of R (0.08201 instead of 0.0821 or 0.08206) does not affect the calculated value of n when the correct kelvin temperature is used and the answer is reported to three significant figures.

Question 5 (continued)**Sample: 5C****Score: 1****Point 01: 0**

Part A: The point was not earned because the response does not identify the geometry around the Si atom as tetrahedral.

Point 02: 0

Part B: The point was not earned. While the response correctly agrees with the claim, it does not provide a justification.

Point 03: 1

Part C: The point was earned for correctly identifying that compound X has the greater vapor pressure because it has reached its boiling point, while compound Y has not yet reached its boiling point.

Point 04: 0

Part D: The point was not earned because the response does not demonstrate a correct calculation for the moles of gas in the container. The response adds the molar masses.

