

2017

AP[®]

CollegeBoard

AP Biology

Free-Response Questions

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

BIOLOGY

Section II

8 Questions

Total Time—90 minutes

Reading Period—10 minutes

Writing Period—80 minutes

Directions: Questions 1 and 2 are long free-response questions that require about 22 minutes each to answer and are worth 10 points each. Questions 3–8 are short free-response questions that require about 6 minutes each to answer. Questions 3–5 are worth 4 points each and questions 6–8 are worth 3 points each.

Read each question carefully and completely. You are advised to spend the 10-minute reading period planning your answers. You may begin writing your responses before the reading period is over. Write your response in the space provided for each question. Only material written in the space provided will be scored. Answers must be written out in paragraph form. Outlines, bulleted lists, or diagrams alone are not acceptable.

TABLE 1. EFFECT OF 0.1 mM CAFFEINE ON MEMORY IN BEES


Treatment	Memory (average probability of revisiting a nectar source $\pm 2SE_{\bar{x}}$)	
	10 Minutes	24 Hours
Control	0.72 ± 0.09	0.41 ± 0.07
Caffeine	0.83 ± 0.07	0.78 ± 0.08

1. In flowering plants, pollination is a process that leads to the fertilization of an egg and the production of seeds. Some flowers attract pollinators, such as bees, using visual and chemical cues. When a bee visits a flower, in addition to transferring pollen, the bee can take nectar from the flower and use it to make honey for the colony.

Nectar contains sugar, but certain plants also produce caffeine in the nectar. Caffeine is a bitter-tasting compound that can be toxic to insects at high concentrations. To investigate the role of caffeine in nectar, a group of researchers studied the effect of 0.1 mM caffeine on bee behavior. The results of an experiment to test the effect of caffeine on bees' memory of a nectar source are shown in Table 1.

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

- (a) On the axes provided, **construct** an appropriately labeled graph to illustrate the effect of caffeine on the probability of bees revisiting a nectar source (memory).
- (b) Based on the results, **describe** the effect of caffeine on each of the following:
- Short-term (10 minute) memory of a nectar source
 - Long-term (24 hour) memory of a nectar source
- (c) **Design an experiment** using artificial flowers to investigate potential negative effects of increasing caffeine concentrations in nectar on the number of floral visits by bees. **Identify** the null hypothesis, an appropriate control treatment, and the predicted results that could be used to reject the null hypothesis.
- (d) Researchers found that nectar with caffeine tends to have a lower sugar content than nectar without caffeine. Plants use less energy to produce the caffeine in nectar than they do to produce the sugar in nectar. **Propose ONE benefit** to plants that produce nectar with caffeine and a lower sugar content. **Propose ONE cost** to bees that visit the flowers of plants that produce nectar with caffeine and a lower sugar content.



2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS




Figure 1. The effect of karrikins (KAR) and trimethylbutenolides (TMB) on seed germination in *Lactuca* plants. Error bars represent $\pm 2SE_{\bar{x}}$.

2. Fires frequently occur in some ecosystems and can destroy all above-ground vegetation. Many species of plants in these ecosystems respond to compounds in smoke that regulate seed germination after a major fire. Karrikins (KAR) and trimethylbutenolides (TMB) are water-soluble compounds found in smoke that are deposited in the soil as a result of a fire. KAR and TMB bind to receptor proteins in a seed. In a study on the effects of smoke on seeds, researchers recorded the timing and percent of seed germination in the presence of various combinations of KAR and TMB. The results are shown in Figure 1.

In a second investigation into the effect of available water on seed germination after a fire, researchers treated seeds with KAR or TMB. The treated seeds were then divided into two treatment groups. One group received a water rinse and the other group received no water rinse. The seeds were then incubated along with a group of control seeds that were not treated. The results are shown in the table.

EFFECT OF CHEMICAL TREATMENT AND WATER RINSE ON GERMINATION

Treatment Group	Chemical Treatment		Water	Germination Result
	KAR	TMB		
1 (control)	–	–	–	Control result
2	+	–	–	Different from control
3	–	+	–	Different from control
4 (control)	–	–	+	Control result
5	+	–	+	Different from control
6	–	+	+	Same as control

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

(a) The researchers made the following claims about the effect of KAR and the effect of TMB on seed germination relative to the control treatment.

- KAR alone affects the timing of seed germination.
- KAR alone affects the percentage of seeds that germinate.
- TMB alone affects the timing of seed germination.
- TMB alone affects the percentage of seeds that germinate.

Provide support using data from Figure 1 for each of the researchers' claims.

(b) **Make a claim** about the effect of rinsing on the binding of KAR to the receptor in the seed and about the effect of rinsing on the binding of TMB to the receptor in the seed. Identify the appropriate treatment groups and results from the table that, when compared with the controls, **provide support** for each claim.

(c) There is intense competition by plants to successfully colonize areas that have been recently cleared by a fire. **Describe** ONE advantage of KAR regulation and ONE advantage of TMB regulation to plants that live in an ecosystem with regular fires.

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

3. Gibberellin is the primary plant hormone that promotes stem elongation. GA 3-beta-hydroxylase (GA3H) is the enzyme that catalyzes the reaction that converts a precursor of gibberellin to the active form of gibberellin. A mutation in the *GA3H* gene results in a short plant phenotype. When a pure-breeding tall plant is crossed with a pure-breeding short plant, all offspring in the F₁ generation are tall. When the F₁ plants are crossed with each other, 75 percent of the plants in the F₂ generation are tall and 25 percent of the plants are short.

		Second Base in Codon				
		U	C	A	G	
First Base in Codon	U	UUU } Phe UUC } UUA } Leu UUG }	UCU } UCC } Ser UCA } UCG }	UAU } Tyr UAC } UAA Stop UAG Stop	UGU } Cys UGC } UGA Stop UGG Trp	U C A G
	C	CUU } CUC } Leu CUA } CUG }	CCU } CCC } Pro CCA } CCG }	CAU } His CAC } CAA } Gln CAG }	CGU } CGC } Arg CGA } CGG }	U C A G
	A	AUU } AUC } Ile AUA } AUG Met or Start	ACU } ACC } Thr ACA } ACG }	AAU } Asn AAC } AAA } Lys AAG }	AGU } Ser AGC } AGA } Arg AGG }	U C A G
	G	GUU } GUC } Val GUA } GUG }	GCU } GCC } Ala GCA } GCG }	GAU } Asp GAC } GAA } Glu GAG }	GGU } GGC } Gly GGA } GGG }	U C A G

Figure 1. The universal genetic code


- (a) The wild-type allele encodes a GA3H enzyme with alanine (Ala), a nonpolar amino acid, at position 229. The mutant allele encodes a GA3H enzyme with threonine (Thr), a polar amino acid, at position 229. **Describe** the effect of the mutation on the enzyme and **provide reasoning** to support how this mutation results in a short plant phenotype in homozygous recessive plants.
- (b) Using the codon chart provided, **predict** the change in the codon sequence that resulted in the substitution of alanine for threonine at amino acid position 229.
- (c) **Describe** how individuals with one (heterozygous) or two (homozygous) copies of the wild-type *GA3H* allele can have the same phenotype.

2017 AP® BIOLOGY FREE-RESPONSE QUESTIONS

DIETARY COMPOSITION OF ORGANISMS IN AN AQUATIC ECOSYSTEM

Organism	Food Source (% of diet)				
	Algae	Stoneflies	Midges	Hellgrammites	Caddisflies
Algae					
Stoneflies			90		10
Midges	100				
Hellgrammites		20	10		70
Caddisflies	70		30		

4. The table above shows how much each organism in an aquatic ecosystem relies on various food sources. The rows represent the organisms in the ecosystem, and the columns represent the food source. The percentages indicate the proportional dietary composition of each organism. High percentages indicate strong dependence of an organism on a food source.
- (a) Based on the food sources indicated in the data table, **construct** a food web in the template below. Write the organism names on the appropriate lines AND draw the arrows necessary to indicate the energy flow between organisms in the ecosystem.
- (b) In an effort to control the number of midges, an area within the ecosystem was sprayed with the fungus *Metarhizium anisopliae*, which significantly decreased the midge population. Based on the data in the table, **predict** whether the spraying of the fungus will have the greatest short-term impact on the population of the stoneflies, the caddisflies, or the hellgrammites. **Justify** your prediction.



2017 AP® BIOLOGY FREE-RESPONSE QUESTIONS




Figure 1. Characteristics of a pond community over time

5. *Microcystis aeruginosa* is a freshwater photosynthetic cyanobacterium. When temperatures increase and nutrients are readily available in its pond habitat, *M. aeruginosa* undergoes rapid cell division and forms an extremely large, visible mass of cells called an algal bloom. *M. aeruginosa* has a short life span and is decomposed by aerobic bacteria and fungi. **Identify** the metabolic pathway and the organism that is primarily responsible for the change in oxygen level in the pond between times I and II AND between times III and IV.

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS




Figure 1. Comet assay to detect double-stranded breaks in DNA

6. A comet assay is a technique used to determine the amount of double-strand breaks in DNA (DNA damage) in cells. The nucleus of an individual cell is placed on a microscope slide coated with an agarose gel. An electric current is applied to the gel that causes DNA to move (electrophoresis), and the DNA is stained with a fluorescent dye. When viewed using a microscope, undamaged DNA from the nucleus appears as a round shape (the head), and the fragments of damaged DNA extend out from the head (the tail). The length of the tail corresponds to the amount of the damage in the DNA (see Figure 1).
- (a) To explain the movement of DNA fragments in the comet assay, **identify** one property of DNA and **provide reasoning** to support how the property contributes to the movement during the comet assay technique.
- (b) In a different experiment, cells are treated with a chemical mutagen that causes only nucleotide substitutions in DNA. **Predict** the likely results of a comet assay for this treatment.

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

7. Many species of bacteria grow in the mouths of animals and can form biofilms on teeth (plaque). Within plaque, the outer layers contain high levels of oxygen and the layers closest to the tooth contain low levels of oxygen. The surface of the tooth is covered in a hard layer of enamel, which can be dissolved under acidic conditions. When the enamel breaks down, the bacteria in plaque can extract nutrients from the tooth and cause cavities.

Certain types of bacteria (e.g., *Streptococcus mutans*) thrive in the innermost anaerobic layers of the plaque and are associated with cavities. Other types of bacteria (*Streptococcus sanguinis*) compete with *S. mutans* but are unable to thrive in acidic environments.

- (a) **Identify** the biochemical pathway *S. mutans* uses for metabolizing sugar and **describe** how the pathway contributes to the low pH in the inner layers of plaque.
- (b) Normal tooth brushing effectively removes much of the plaque from the flat surfaces of teeth but cannot reach the surfaces between teeth. Many commercial toothpastes contain alkaline components, which raise the pH of the mouth. **Predict** how the population sizes of *S. mutans* AND *S. sanguinis* in the bacterial community in the plaque between the teeth are likely to change when these toothpastes are used.

2017 AP[®] BIOLOGY FREE-RESPONSE QUESTIONS

8. Estrogens are small hydrophobic lipid hormones that promote cell division and the development of reproductive structures in mammals. Estrogens passively diffuse across the plasma membrane and bind to their receptor proteins in the cytoplasm of target cells.
- (a) **Describe** ONE characteristic of the plasma membrane that allows estrogens to passively cross the membrane.
 - (b) In a laboratory experiment, a researcher generates antibodies that bind to purified estrogen receptors extracted from cells. The researcher uses the antibodies in an attempt to treat estrogen-dependent cancers but finds that the treatment is ineffective. **Explain** the ineffectiveness of the antibodies for treating estrogen-dependent cancers.

STOP

END OF EXAM